

# The Sources of Inflammatory Mediators in the Lung Following Silica Exposure

**K. Murali Krishna Rao, Dale W. Porter, Terence Meighan,  
and Vince Castranova**

**doi:10.1289/ehp.7295 (available at <http://dx.doi.org/>)**

**Online 16 August 2004**



**The Sources of Inflammatory Mediators in the  
Lung Following Silica Exposure**

**K. Murali Krishna Rao, Dale W. Porter, Terence Meighan, and Vince Castranova**

**Pathology and Physiology Research Branch  
Health Effects Laboratory Division  
National Institute for Occupational Safety and Health  
Morgantown, WV 26505**

**Correspondence: K. Murali Krishna Rao, M.D.  
Box 2015  
PPRB/HELD/NIOSH  
1095 Willowdale Road  
Morgantown, WV 26505  
Tel: 304-285-6169  
FAX: 304-285-5938  
e-mail: [mir8@cdc.gov](mailto:mir8@cdc.gov)**

**Running Title:** Silica-induced inflammatory mediators

**Key Words:** Alveolar Macrophages, Alveolar type II cells, Cytokines, Fibroblasts, Lung, Silica, Gene Expression

**Abbreviations:**

AM - alveolar macrophage, BAL - bronchoalveolar lavage, ELISA - enzyme-linked immunoassays, GM-CSF - granulocyte/macrophage-colony stimulating factor, iNOS - inducible nitric oxide synthase, ICAM-1 - intercellular adhesion molecule-1, IL – interleukin, MIP-2 - macrophage inflammatory protein-2, MCP-1 - monocyte chemoattractant protein-1, PMN – polymorphonuclear neutrophils, TGF -transforming growth factor, TNF- $\alpha$  - tumor necrosis factor- $\alpha$ , type II cells - alveolar epithelial type II cells.

1. Abstract
2. Introduction
3. Materials and Methods
4. Results
5. Discussion
6. References
7. Figure Legends
8. Tables
9. Figures

### Abstract

The expression of 10 genes implicated in regulation of the inflammatory processes in the lung was studied following exposure of alveolar macrophages (AM) to silica *in vitro* or *in vivo*. Exposure of AM to silica *in vitro* upregulated the mRNA levels of three genes (IL-6, MCP-1, and MIP-2) without a concomitant increase in the protein levels. AM isolated following intratracheal instillation of silica upregulated mRNA levels of four additional genes (GM-CSF, IL-1 $\beta$ , IL-10 and iNOS). Protein levels IL-6, MCP-1 and MIP-2 were elevated in bronchoalveolar lavage (BAL) fluid. Fibroblasts under basal culture conditions express much higher levels of IL-6 and GM-CSF compared to AM. Co-culture of AM and alveolar type II cells, or co-culture of AM and lung fibroblasts, in contact cultures or Transwell chambers, revealed no synergistic effect. Therefore, such interaction does not explain the effects seen *in vivo*. In conclusion, identification of the intercellular communication *in vivo* is still unresolved. However, fibroblasts appear to be an important source of inflammatory mediators in the lung.